arXiv:0909.3179v1 [math.SG] 17 Sep 2009

THE FUNDAMENTAL GROUP OF G-MANIFOLDS

HUI LI

Abstract. Let M be a connected smooth G-manifold, where G is a connected compact Lie group. In this paper, we first study the relation between $\pi_1(M)$ and $\pi_1(M/G)$. Then we particularly focus on the case when M is a connected Hamiltonian G-manifold with an equivariant moment map ϕ. In [13], for *compact* M, we proved that $\pi_1(M) \cong \pi_1(M/G) \cong \pi_1(M_a)$ for all $a \in \text{image}(\phi)$, where M_a is the symplectic quotient at a. We generalize and extend these results to Hamiltonian G-manifolds with *proper* moment maps. Our main results are Proposition 1, Theorems 2 and 3.

1. Introduction

Let M be a smooth manifold. Let a connected compact Lie group G act on M. We call M a *G-manifold*.

Let (M, ω) be a symplectic manifold. Assume that a connected compact Lie group G acts on M with moment map $\phi : M \to \mathfrak{g}^*$, where $\mathfrak{g}^*$ is the dual Lie algebra of G. In this case, we call (M, ω) a *Hamiltonian G-manifold*. We will always assume that ϕ is equivariant with respect to the G action, where G acts on $\mathfrak{g}^*$ by the coadjoint action. Take a moment map value a in $\mathfrak{g}^*$, the space $\mathbf{M_a} = \phi^{-1}(G \cdot a)/G$ is called the **symplectic quotient** or the **reduced space** at the coadjoint orbit $G \cdot a$. If G_a is the stabilizer group of a under the coadjoint action, by equivariance of the moment map, $\mathbf{M_a} = \phi^{-1}(a)/G_a = \phi^{-1}(G \cdot a)/G$. We will use the two notations interchangeably.

In this paper, fundamental groups always mean the fundamental groups of topological spaces, even when we do not explicitly say so.

For compact Hamiltonian G-manifold M, in [13], we had the following results.

Theorem 1. *Let (M, ω) be a connected compact Hamiltonian G-manifold with equivariant moment map ϕ, where G is a connected compact Lie group. Then, as fundamental groups of topological spaces, $\pi_1(M) \cong \pi_1(M/G) \cong \pi_1(M_a)$ for all $a \in im(\phi)$.*

If M is not compact, the results in Theorem 1 may not be true as the following easy example shows.

Key words and phrases. Symplectic manifold, fundamental group, Hamiltonian group action, moment map, symplectic quotient.

2000 MSC. Primary : 53D05, 53D20; Secondary : 55Q05, 57R19.

Example 1.1. Consider $M = S^1 \times \mathbb{R}$, and let S^1 rotate the first factor, then the moment map is the projection to the second factor $\mathbb{R}$. We have that $\pi_1\left(M/S^1\right) \cong \pi_1\left(M_a\right) \ncong \pi_1\left(M\right)$ for all $a \in \mathrm{im}(\phi)$.

The proof of Theorem 1 goes as follows (see [13] and [12]): we prove that $\pi_1\left(M\right) \cong \pi_1\left(M_a\right)$ for *some value* a which is a *boundary vertex* of the moment polytope. In fact, for this special value a, the whole set $\phi^{-1}(a)$ is fixed by $G_a \supseteq T$, where G_a is the stabilizer group of a under the coadjoint action, and T is a maximal torus of G. Therefore, $\phi^{-1}(a) \approx \phi^{-1}(a)/G_a = M_a$. So

$$\pi_1\left(M\right) \cong \pi_1\left(\phi^{-1}(a)\right) \cong \pi_1\left(M_a\right). \tag{1.2}$$

Then we prove that the $\pi_1(M_a)$'s are isomorphic for all value a's. Finally, we prove that $\pi_1\left(M/G\right) \cong \pi_1\left(M_a\right)$ for a particular value a.

Lemma 1.3. *Let (M,ω) be a connected Hamiltonian G-manifold, where G is a connected compact Lie group. Let T be a maximal torus of G. If M is* compact*, then there exists a connected subgroup H of G such that $H \supseteq T$ and that H has a fixed point.*

Proof. Let ϕ be the moment map. If $G = T$ is abelian, then $\phi(M)$ is a convex polytope for compact M. The inverse image of each boundary vertex of the polytope is a fixed component of T. If G is nonabelian, let a be a furthest value in the image of ϕ. In Lemma 6.14 of [13], although we did not state it, we proved that $\phi^{-1}(a)$ is fixed by $G_a \supseteq T$, where G_a is the stabilizer group of a under the coadjoint action. □

If M is not compact, as we saw in Example 1.1, there may not be isotropy group which contains a maximal torus of G; and, for this example, $\pi_1\left(M\right) \ncong \pi_1\left(M/G\right)$. While for compact M, Lemma 1.3 holds, and Theorem 1 says that $\pi_1\left(M\right) \cong \pi_1\left(M/G\right)$.

Now, we only assume that M is a connected G-manifold with moment map ϕ. The moment map ϕ is **proper** if the inverse image under ϕ of each compact set is compact.

For a noncompact Hamiltonian G-manifold M with proper moment map ϕ, $\phi(M) \cap \mathfrak{t}^*_+$ is a convex polyhedral set (Theorem 4), where $\mathfrak{t}^*_+$ is a closed positive Weyl chamber. But, $\phi(M)$ may not be compact and hence there may not be boundary vertex on $\phi(M) \cap \mathfrak{t}^*_+$. We will take a different approach from that of compact M to study $\pi_1(M)$. First, we study $\pi_1(M)$ for general G-manifolds.

Let a connected compact Lie group G act on a connected smooth manifold M. Let H be a subgroup of G which is the stabilizer of some point in M. Let (H) be the subgroups of G conjugate to H. Denote $M_{(H)} = \{x \in M \mid G_x \in (H)\}$. The set $M_{(H)}$ is a submanifold of M, and we call it the **(H)-orbit type** or the **(H)-isotropy type**. There is a *connected open dense* orbit type in M, which is called the **principal orbit type**. We first have the following result.

Proposition 1. *Let G act on a connected smooth manifold M, where G is a connected compact Lie group. Assume that each nonprincipal orbit type has codimension at least 2, and that there exists an orbit which is simply connected. Then*

$$\pi_1(M) \cong \pi_1\left(M/G\right).$$

If we assume that big enough isotropy group $H \supseteq T$ exists for the G action, then there exists a simply connected orbit (see Lemma 4.2). Now, let us consider Hamiltonian G-manifolds. Our next main result is Theorem 2, where (A) holds without assuming the moment map is proper, and we assume the moment map is proper in (B).

Theorem 2. *Let (M, ω) be a connected Hamiltonian G-manifold with equivariant moment map ϕ, where G is a connected compact Lie group. Let T be a maximal torus of G. Assume that there is a connected subgroup H of G such that $H \supseteq T$ and that H has a fixed point. Then*

(A) $\pi_1\left(M\right) \cong \pi_1\left(M/G\right) \cong \pi_1\left(M/T\right) \cong \pi_1\left(M/H\right)$.

(B) *Let x be the H fixed point. If ϕ is proper, and if $\phi(x) = a$, then*

$$\pi_1\left(\phi^{-1}(G\cdot a)\right) \cong \pi_1\left(\phi^{-1}(a)\right) \cong \pi_1\left(M_a\right).$$

Remark 1.4. In Theorem 2, if there is a subgroup H such that $H \supseteq T$ has a fixed point, then the identity component of H has a fixed point. So we can always take connected H. In fact, all the isomorphisms except $\pi_1\left(M\right) \cong \pi_1\left(M/H\right)$ do not need H to be connected. We take a connected group to make M to be a Hamiltonian H-manifold for the connected group H.

Next, we observe that, the fact $\pi_1\left(M/G\right) \cong \pi_1\left(M_a\right)$ for all $a \in \mathrm{im}(\phi)$ always holds for Hamiltonian G-manifolds with proper equivariant moment maps. This is a straightforward generalization of the compact case. See Section 5 for the proof. We observe that this result does not depend on whether the action has big enough isotropy groups. In our next main result Theorem 3 below, we first state this generalization in (A). In (B), we made a step further in generalizing (A). For noncompact M, we observed that there may not be isotropy groups which contain a maximal torus of G. If we take $\widehat{G}$ to be a connected subgroup of G such that up to finitely many elements, every isotropy group is a subgroup of $\widehat{G}$, we can express $\pi_1\left(M/\widehat{G}\right)$ in terms of π_1 of the "partial" reductions of the G-moment map by $\widehat{G}$.

Theorem 3. *Let (M, ω) be a connected Hamiltonian G-manifold with a proper equivariant moment map ϕ, where G is a connected compact Lie group.*

(A) $\pi_1\left(M/G\right) \cong \pi_1\left(M_a\right)$ *for all* $a \in im(\phi)$.

(B) *More generally, let $\widehat{G}$ be any connected subgroup of G such that up to finitely many elements, every isotropy group is a subgroup of $\widehat{G}$. Then,* $\pi_1\left(M/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(G\cdot a)/\widehat{G}\right)$ *for all* $a \in im(\phi)$.

A special case is when $\mathrm{im}(\phi)$ consists of regular values. In this case, all the isotropy groups are finite. Taking $\widehat{G} = 1$ in Theorem 3, we obtain $\pi_1(M) \cong \pi_1(\phi^{-1}(G\cdot a))$ for all $a \in \mathrm{im}(\phi)$. We have the following lemma, where the proof of the fact $\pi_1(M) \cong \pi_1\left(\phi^{-1}(a)\right)$ for all $a \in \mathrm{im}(\phi)$ is given below.

Lemma 1.5. *Let (M,ω) be a connected Hamiltonian G-manifold with a proper equivariant moment map ϕ, where G is a connected compact Lie group. Assume that image(ϕ) consists of regular values. Then,*

$$\pi_1(M) \cong \pi_1\left(\phi^{-1}(a)\right) \cong \pi_1\left(\phi^{-1}(G\cdot a)\right) \quad \textit{for all} \quad a \in \mathit{im}(\phi).$$

Proof of $\pi_1(M) \cong \pi_1\left(\phi^{-1}(a)\right)$ for all $a \in im(\phi)$. If $\mathrm{im}(\phi)$ consists of regular values, then ϕ is a proper submersion from M onto $\mathrm{im}(\phi) \subset \mathfrak{g}^*$. By Ehresmann's lemma, ϕ gives a fibration from M onto $\mathrm{im}(\phi)$ (which is open in $\mathfrak{g}^*$) with connected fiber ([8] or [11]) diffeomorphic to $\phi^{-1}(a)$ for *any* $a \in \mathrm{im}(\phi)$. By the homotopy exact sequence for fibrations, we have $\pi_1(M) \cong \pi_1\left(\phi^{-1}(a)\right)$. □

We have the following corollary of Theorems 2 and 3. It states the important contribution to $\pi_1(M)$ of the level sets which contain fixed points of big enough subgroups of G. In particular, this is a generalization of the fact $\pi_1(M) \cong \pi_1(\text{minimum}) \cong \pi_1(\text{maximum})$ for a compact Hamiltonian S^1-manifold M, where minimum and maximum are those of the real valued moment map for the circle action ([12]); and, more generally, it is a generalization of $\pi_1(M) \cong \pi_1\left(\phi^{-1}(a)\right)$ as in (1.2) for some boundary vertices of the moment polytope for compact Hamiltonian G-manifold M. This corollary and (B) of Theorem 2 together generalize (1.2), even though here H may not fix the whole set $\phi^{-1}(a)$.

Corollary 1.6. *Let (M,ω) be a connected Hamiltonian G-manifold with a proper equivariant moment map ϕ, where G is a connected compact Lie group. Let T be a maximal torus of G. Assume that there is a connected subgroup H of G such that $H \supseteq T$ and that H has a fixed point x, and $\phi(x) = a$. Then*

$$\pi_1(M) \cong \pi_1\left(\phi^{-1}(G\cdot a)\right) \cong \pi_1\left(\phi^{-1}(a)\right).$$

For S^1-manifolds with proper moment maps, in [7], using Morse theory, Godinho and Sousa-Dias proved that when S^1 does not have fixed points, $\pi_1(M) \cong \pi_1\left(\phi^{-1}(a)\right)$ for all $a \in \mathrm{im}(\phi)$; and, when S^1 has a fixed point, $\pi_1(M) \cong \pi_1(M_a)$ for all $a \in \mathrm{im}(\phi)$. This result is a partial result of our Theorems 2 and 3 for the case $G = S^1$.

2. PRELIMINARIES ON G-MANIFOLDS AND ON HAMILTONIAN G-MANIFOLDS

In this section, we review some theorems on Hamiltonian G-manifolds which constitute the main tools in our proofs, and we will explain how we use them. We review the notion of stratified spaces which are essential for our proof of Theorem 3 since M/G and M_a are in general stratified spaces.

Theorem 4. *([1], [4], [8] and [11]) Let (M, ω) be a connected Hamiltonian G-manifold with proper moment map ϕ, where G is a connected compact Lie group. Let $\mathfrak{t}^*_+$ be a fixed closed positive Weyl chamber of $\mathfrak{g}^*$. Then*

1. *For each coadjoint orbit $\mathcal{O}$ in the image of ϕ, $\phi^{-1}(\mathcal{O})$ is connected.*
2. *The image $\phi(M) \cap \mathfrak{t}^*_+$ is a convex polyhedral set.*

If $G = T$ is abelian, then $\phi(M)$ is a convex polyhedral set; it consists of **faces** with different dimensions. A connected open face is called a **chamber**. By quotienting out a subtorus which acts trivially, we can assume that the action is effective, and hence there is an open face.

Theorem 5. *([16] or [10]) Let (M, ω) be a connected Hamiltonian G-manifold with proper moment map ϕ, where G is a connected compact Lie group. Choose a G-invariant metric on M. Assume that $0 \in im(\phi)$. Then there exists a G-invariant open neighborhood $U \subset \mathfrak{g}^*$ of 0 such that the gradient flow of the function $-\|\phi\|^2$ induces a G-equivariant deformation retraction from $\phi^{-1}(U)$ to $\phi^{-1}(0)$.*

2.1. Local normal form theorem.

Theorem 6. *(Local normal form) ([14], [6]) Let (M, ω) be a symplectic manifold with a Hamiltonian connected compact Lie group G action. Let $p \in M$, and assume that the orbit $G \cdot p$ is isotropic. Let H be the isotropy subgroup of p. Then a neighborhood of the orbit $G \cdot p$ in M is equivariantly symplectomorphic to $G \times_H (\mathfrak{h}^\circ \times V)$, where $\mathfrak{h}^\circ$ is the annihilator of $\mathfrak{h}$ =Lie(H) in $\mathfrak{g}^*$ on which H acts by the coadjoint action, and V is a complex vector space on which H acts linearly and symplectically.*

The G action on this local model is $g_1 \cdot [g, a, v] = [g_1 g, a, v]$, and the moment map on this local model is $\phi([g, a, v]) = Ad^*(g)(\phi(p) + a + \psi(v))$, where $\psi(v)$ is the moment map for the H action on V.

2.2. Cross section theorem.

Definition 2.1. Suppose that a group G acts on a manifold M. Given a point m in M with isotropy group G_m, a submanifold $U_m \subset M$ containing m is a **slice at m** if U_m is G_m-invariant, $G \cdot U_m$ is a neighborhood of m, and the map

$$G \times_{G_m} U_m \to G \cdot U_m, \qquad [g, u] \longmapsto g \cdot u \quad \text{is an isomorphism.}$$

For instance, consider the co-adjoint action of $G = SU(2)$ or $SO(3)$ on $\mathbb{R}^3$ =Lie(G). For $a \in \mathbb{R}^3, a \neq 0$, there is a unique ray I_a passing through 0 and a; the open ray $I_a^\circ = I_a - 0$ is a slice at a. If $a = 0$, then a slice at 0 is $\mathbb{R}^3$.

More generally, consider the co-adjoint action of a connected compact Lie group G on $\mathfrak{g}^*$. Fix a closed positive Weyl chamber $\mathfrak{t}^*_+$, without loss of generality, take $a \in \mathfrak{t}^*_+$. Let $\tau \subset \mathfrak{t}^*_+$ be the open face of $\mathfrak{t}^*_+$ containing a and let G_a be the isotropy group of a. Since all the points on τ have the same isotropy group, we also use G_τ to denote G_a. Then the natural slice at a is $U_a = G_a \cdot \{b \in \mathfrak{t}^*_+ \mid G_b \subset G_a\} = G_a \cdot \bigcup_{\tau \subset \bar{\tau'}} \tau'$, and it is an open subset of $\mathfrak{g}^*_\tau = \mathfrak{g}^*_a$.

The following cross section theorem is due to Guillemin and Sternberg. (Theorem 26.7 in [5]; for the following version, see Corollary 2.3.6 in [3].)

Theorem 7. *(cross section). Let (M, ω) be a symplectic manifold with a moment map $\phi : M \to \mathfrak{g}^*$ arising from an action of a compact connected Lie group G. Let a be a point in $\mathfrak{g}^*$ and let U_a be the natural slice at a. Then the* **cross section** *$R = \phi^{-1}(U_a)$ is a G_a-invariant symplectic submanifold of M, where G_a is the isotropy group of a. Furthermore, the restriction $\phi|_R$ is a moment map for the action of G_a on R.*

The highest dimensional face τ^P of $\mathfrak{t}^*_+$ which contains values of ϕ is called the **principal face**. Let U_{τ^P} be the slice at τ^P. The cross section $\phi^{-1}(U_{\tau^P})$ is called the **principal cross section**. The maximal torus of G acts on the principal cross section (but not necessarily effectively, see Theorem 3.1 in [11]).

We will use the cross section theorem in the following two ways:

- If $a \in \text{im}(\phi) \cap \tau$ for some face $\tau \subset \mathfrak{t}^*_+$, then τ lies on the central dual Lie algebra of G_a, so the G_a-orbits in $\phi^{-1}(a)$ are isotropic orbits; then we can use the local normal form theorem in the cross section R to describe a neighborhood of a G_a-orbit.
- Assume that the moment map ϕ is proper. Fix a G-invariant Riemannian metric on M. Assume that $a \in \text{im}(\phi) \cap \tau$. By Theorem 5, there exists a G_a-invariant neighborhood of $\phi^{-1}(a)$ in R which G_a-equivariantly deformation retracts to $\phi^{-1}(a)$. By equivariance of ϕ, there exists a G-invariant neighborhood $\mathcal{N}$ of $\phi^{-1}(G \cdot a)$ in M which G-equivariantly deformation retracts to $\phi^{-1}(G \cdot a)$.

2.3. Stratified space and link of a point.

Let G, a connected compact Lie group, act on a smooth manifold M, the quotient space M/G is a stratified space. If (M, ω) is a symplectic manifold and the G action is Hamiltonian, then the symplectic quotients are stratified spaces ([15]).

Let X be a Hausdorff and paracompact topological space and let $\mathcal{J}$ be a partially ordered set with order relation denoted by $\leq$. A $\mathcal{J}$-decomposition of X is a locally finite collection of disjoint, locally closed manifolds $S_i \subset X$ (one for each $i \in \mathcal{J}$) called pieces such that $X = \cup_{i \in \mathcal{J}} S_i$, and that $S_i \cap \bar{S}_j \neq \emptyset \Leftrightarrow S_i \subset \bar{S}_j \Leftrightarrow i \leq j$. We call the space X a $\mathcal{J}$- **decomposed space**.

A stratified space is defined recursively as follows. A decomposed space X is called a **stratified space** if the pieces of X, called strata, satisfy the following condition: Given a point x in a piece S (connected), there exists an open neighborhood $\tilde{U}$ of x in X, an open ball B around x in S, a compact stratified space L, called the **link of x**, and a homeomorphism

$$\varphi : B \times \overset{\circ}{C}L \to \tilde{U}$$

that preserves the decompositions. Here, $\overset{\circ}{C}L$ is the space obtained by collapsing the boundary $L \times 0$ of the half-open cylinder $L \times [0, \infty)$ to a point. We also call the link of x the **link of S**.

For a smooth manifold M, if N is a closed submanifold of M, by the tubular neighborhood theorem, a neighborhood of N in M is diffeomorphic to a neighborhood of the normal bundle E of N in M. If the fiber sphere of the sphere bundle $S(E)$ of N is connected and simply connected, i.e., if the codimension of N is bigger than or equal to 3, then $\pi_1(M) \cong \pi_1(M - N)$. For a stratified space X, the *link* of a stratum S plays the role of the fiber sphere of the sphere bundle of a submanifold in a smooth manifold. Using the Van-Kampen theorem, we can prove the following lemma. May see Lemma 2.3 in [13] for detail.

Lemma 2.2. *Let X be a stratified space. Let S be a connected closed stratum of X whose link is connected and simply connected. Then $\pi_1(X) \cong \pi_1(X - S)$.*

3. Proof of Proposition 1

Lemma 3.1. *(Corollary 6.3 in [2]) If X is an arcwise connected G-space, G compact Lie, and if there is an orbit which is connected, then the fundamental group of X maps onto that of X/G.*

Let M be a connected smooth G-manifold, where G is a connected compact Lie group. In the Introduction, we defined the (H)-orbit type $M_{(H)}$ for a stabilizer group H and we defined the principal orbit type. The fact is that all other orbit types are in the closure of the principal orbit type. More generally, if $M_{(H)} \subset \bar{M}_{(H')}$, then $(H') \subset (H)$. The quotient M/G is a stratified space whose strata are the $M_{(H)}/G$'s. The manifold $M_{(H)}/G$ is called the **(H)-stratum** of M/G.

Lemma 3.2. *Let G act on a connected smooth manifold M, where G is a connected compact Lie group. Let $M_{(H)}$ be a closed connected orbit type which has codimension at least 2 in M. Then we have a surjection*

$$\pi_1\left(M - M_{(H)}\right) \twoheadrightarrow \pi_1(M).$$

Proof. Since the submanifold $M_{(H)}$ has codimension at least 2 in M, its sphere bundle $S\left(M_{(H)}\right)$ (in particular the fiber) is connected, by the homotopy exact sequence for the fibration $S\left(M_{(H)}\right) \to M_{(H)}$, we have a surjection

$$\pi_1\left(S\left(M_{(H)}\right)\right) \twoheadrightarrow \pi_1\left(M_{(H)}\right).$$

Take the open cover $\{M - M_{(H)}, N\left(M_{(H)}\right)\}$ of M, where $N\left(M_{(H)}\right)$ is a small open neighborhood of $M_{(H)}$. By the Van-Kampen theorem, we have a surjection

$$\pi_1\left(M - M_{(H)}\right) \twoheadrightarrow \pi_1(M).$$

$\square$

Remark 3.3. Let G act on a connected smooth manifold M, where G is a connected compact Lie group. Assume that *each* nonprincipal orbit type has codimension at least 2 in M. Using Lemma 3.2 inductively, in a suitable order, we can continue to throw away certain orbit types to get a surjection. For example, if $\mathcal{M}_{(H)}$ is the submanifold which consists of those orbit types whose closures contain $M_{(H)}$, where $M_{(H)}$ is a closed connected orbit type, then

$$\pi_1\left(\mathcal{M}_{(H)}\right) \twoheadrightarrow \pi_1(M);$$

and, if M_P is the principal orbit type, then we have a surjection

$$\pi_1(M_P) \twoheadrightarrow \pi_1(M).$$

Proof of Proposition 1. Since G is connected, by Lemma 3.1, we have a surjection

$$\pi_1(M) \twoheadrightarrow \pi_1(M/G).$$

Let $G \cdot x$ be the simply connected orbit. Let $q\colon M \to M/G$ be the quotient map and let $\bar{x} = q(x)$. Let α be a loop in M/G based at $\bar{x}$ such that $[\alpha] = 1 \in \pi_1(M/G; \bar{x})$. Let $\widetilde{\alpha}$ be a loop in M based at x which is any lift of α. We need to show that $[\widetilde{\alpha}] = 1 \in \pi_1(M; x)$.

First note that in each orbit type, the orbits are diffeomorphic to each other, so we can use this diffeomorphism to deform paths or loops in the same orbit type. Secondly, for a nonprincipal orbit type, since its codimension in M is at least 2, by Lemma 3.2, we can push paths or loops out of this nonprincipal orbit type to its sphere bundle. For the loop $\widetilde{\alpha}$, we need to deform it through a family of base point preserving loops to the constant loop x. Since $[\alpha] = 1 \in \pi_1(M/G; \bar{x})$, we have a continuous family of paths from each point of α to $\bar{x}$. In M, we follow this continuous family of paths between the orbits which intersect $\widetilde{\alpha}$ and the orbit $G \cdot x$, and we use the above two types of deformation to deform $\widetilde{\alpha}$ to a loop $\widetilde{\alpha}'$ which lies in the orbit $G \cdot x$. Since $\pi_1(G \cdot x) = 1$, $[\widetilde{\alpha}] = [\widetilde{\alpha}'] = 1 \in \pi_1(M; x)$. $\square$

Remark 3.4. Using Proposition 1 on a subspace $\mathcal{M}_{(H)}$ as follows, we have

Lemma 3.5. *Let G, a connected compact Lie group, act on a connected smooth manifold M. Let $M_{(H)}$ be a connected orbit type whose orbits are simply connected. Let $\mathcal{M}_{(H)}$ be the submanifold which consists of those orbit types whose closures contain $M_{(H)}$. Assume that each nonprincipal orbit type in $\mathcal{M}_{(H)}$ has codimension at least 2 in $\mathcal{M}_{(H)}$. Then*

$$\pi_1\left(\mathcal{M}_{(H)}\right) \cong \pi_1\left(\mathcal{M}_{(H)}/G\right).$$

4. Proof of Theorem 2

Lemma 4.1. *Let M be a connected Hamiltonian G-manifold, where G is a connected compact Lie group. Then, each nonprincipal orbit type has codimension at least* 2.

Proof. First assume that $G = T$ is abelian. By the local normal form theorem, a neighborhood of an orbit with isotropy group H is isomorphic to $A = T \times_H (\mathfrak{h}^\circ \times V) = T \times_H (\mathfrak{h}^\circ \times W \times V^H)$, where V^H is the subspace of V fixed by H and W is its complementary subspace in V. Note that $A_H = T \times_H (\mathfrak{h}^\circ \times V^H)$. So if H is not the principal stabilizer, then W is a nontrivial complex vector space on which H acts nontrivially; so M_H has codimension at least 2.

Now assume that G is nonabelian. Fix an isotropy type (H). We may assume that the image of the orbit type $M_{(H)}$ intersects a face τ of $\mathfrak{t}^*_+$ (then $H \subset G_\tau$). By equivariance of the moment map, we only need to consider this orbit type in the cross section R on which G_τ acts. By the local normal form theorem, a neighborhood of an isotropic orbit in R with stabilizer (H) is isomorphic to $A = G_\tau \times_H (\mathfrak{h}^\circ \times V) = G_\tau \times_H (\mathfrak{h}^\circ \times W \times V^H)$, where $\mathfrak{h}^\circ$ is the annihilator of $\mathfrak{h}$ =Lie(H) in $\mathfrak{g}^*_\tau$. Then $A_{(H)} = G_\tau \times_H (\mathbb{R}^m \times V^H)$, where $\mathbb{R}^m$ is a subspace of $\mathfrak{h}^\circ$ fixed by H. Let $\mathbb{R}^l$ be the complementary subspace of $\mathbb{R}^m$ in $\mathfrak{h}^\circ$. If (H) is not the principal stabilizer type, then $\mathbb{R}^l \oplus W \neq \emptyset$. Based on Proposition 6.3 in [13] (one may refer to the proof of Lemma 6.19 in [13] for all possible H), $\dim(\mathbb{R}^l \oplus W) \geq 2$. □

Lemma 4.2. *Let G be a connected compact Lie group, and let T be a maximal torus of G. Let H be a subgroup of G such that $T \subseteq H$. Then $\pi_1(G/H) = 1$.*

Proof. Since we have a surjection

$$\pi_1(T) \twoheadrightarrow \pi_1(G) \quad \text{and the inclusions}$$

$$T \overset{\iota}{\hookrightarrow} H \overset{\iota}{\hookrightarrow} G,$$

we have a surjection

$$\pi_1(H) \twoheadrightarrow \pi_1(G).$$

The claim follows from the homotopy exact sequence for the fibration $H \subset G \to G/H$:

$$\cdots \to \pi_1(H) \to \pi_1(G) \to \pi_1(G/H) \to \cdots .$$

□

Proposition 2. *Let M be a connected Hamiltonian G-manifold with proper equivariant moment map ϕ, where G is a connected compact Lie group. Let T be a maximal torus of G. Assume that there is a subgroup H of G such that $H \supseteq T$ and that H has a fixed point x, and $\phi(x) = a$. Then*

$$\pi_1\left(\phi^{-1}(G \cdot a)\right) \cong \pi_1\left(\phi^{-1}(a)\right) \cong \pi_1(M_a).$$

Proof. Since ϕ is proper, $\phi^{-1}(G\cdot a)$ and $\phi^{-1}(a)$ are connected by Theorem 4; and, by definition of properness, $\phi^{-1}(G\cdot a)$ and $\phi^{-1}(a)$ are also compact. Hence, M_a is connected and compact.

Consider the map

$$\pi_1\left(\phi^{-1}(G\cdot a)\right) \to \pi_1\left(M_a\right). \tag{4.3}$$

Since G is connected and it acts on $\phi^{-1}(G\cdot a)$, by Lemma 3.1, (4.3) is onto.

Now we show that (4.3) is injective. Let $\mathcal{N}$ be an open invariant neighborhood of $\phi^{-1}(G\cdot a)$ in M which equivariantly deformation retracts to $\phi^{-1}(G\cdot a)$. Since the orbit $G\cdot x \approx G/H$, by Lemma 4.2, $\pi_1\left(G\cdot x\right) = 1$. By Lemma 4.1, each nonprincipal orbit type in $\mathcal{N}$ has codimension at least 2. Take $[\alpha] = 1 \in \pi_1\left(M_a\right)$. Then $[\alpha] = 1 \in \pi_1\left(\mathcal{N}/G\right)$. Let $\widetilde{\alpha} \subset \phi^{-1}(G\cdot a)$ be a lifted loop of α. By Proposition 1, $[\widetilde{\alpha}] = 1 \in \pi_1\left(\mathcal{N}\right)$. Since $\pi_1\left(\mathcal{N}\right) = \pi_1\left(\phi^{-1}(G\cdot a)\right)$, $[\widetilde{\alpha}] = 1 \in \pi_1\left(\phi^{-1}(G\cdot a)\right)$.

Now, we prove that $\pi_1\left(\phi^{-1}(a)\right) \cong \pi_1\left(M_a\right)$. Assume $a \in \tau \subset \mathfrak{t}^*_+$. Since H fixes x, and since $\phi(g\cdot x) = Ad^*(g)\cdot\phi(x)$, $H \subset G_a$. Consider the G_a action on the cross section R^τ and on $\phi^{-1}(a)$. Take a G_a-invariant open neighborhood of $\phi^{-1}(a)$ in R^τ which G_a- equivariantly deformation retracts to $\phi^{-1}(a)$ and use the same argument as above. □

Proof of Theorem 2. (A). Let K be G, T or H. Then K is connected and it acts on M. Since M is a Hamiltonian G-manifold, it is a Hamiltonian K-manifold for any K. Moreover, the K action has an orbit which is simply connected by Lemma 4.2. Then $\pi_1\left(M\right) \cong \pi_1\left(M/K\right)$ follows from Lemmas 3.1, 4.1 and Proposition 1.

(B) is Proposition 2. □

5. PROOF OF (A) OF THEOREM 3

The method of removing and deforming used in [13] to prove the same fact for compact manifold M applies to this case. Hamiltonian action plays crucial roles in the removing process — we have the local normal form theorem and the cross section theorem. The *properness of* ϕ or the *compactness of* M is used to control the gradient flow of suitable components of ϕ or of $\|\phi\|^2$ — we use these flows to do *local* deformation retractions (Theorem 5). We will state the important lemmas which lead to the results and we will emphasize the key points which make these lemmas hold. The corresponding proofs in [13] apply or apply with slight modifications.

Lemma 5.1. *(see Lemma 3.6 in* [13]*) Let (M,ω) be a connected Hamiltonian T- manifold with proper moment map ϕ, where T is a connected compact torus. Then, for all value a's in one connected chamber of im(ϕ), $\pi_1(M_a)$'s are isomorphic.*

For the method of removing, the results in Lemmas 5.2 and 5.5 below are the key reasons for Lemmas 7.20, 3.8 and 6.19 in [13] to hold. Lemmas 7.20, 3.8 and 6.19 in [13] were stated in emphasizing the consequences of these results. Here and in Lemmas 6.2 and 7.1, we are particularly emphasizing these points themselves.

Lemma 5.2. *(see Lemmas 7.20 and 3.8 in* [13]*) Let M be a connected Hamiltonian T-manifold with proper moment map ϕ, where T is a connected compact torus. Let $\mathcal{F}$ be a singular face of im(ϕ). Let $\bar{U}$ be the closure of one open connected chamber such that $\mathcal{F} \subset \bar{U}$.*

1. *For each* nonprincipal isotropy type *$M_{(H)} \cap \phi^{-1}(\mathcal{F})$, the link of $\left(M_{(H)} \cap \phi^{-1}(\mathcal{F})\right)/T$ in M/T is connected and simply connected.*
2. *For each isotropy type $M_{(H)} \cap \phi^{-1}(\mathcal{F})$, the link of $\left(M_{(H)} \cap \phi^{-1}(\mathcal{F})\right)/T$ in $\phi^{-1}(\bar{U})/T$ is connected and simply connected.*

We can use (2) of Lemma 5.2 to prove Lemma 5.3. The idea is to use Lemma 2.2 and Theorem 5. We refer to the proof of Lemma 3.8 in [13].

Lemma 5.3. *Let (M, ω) be a connected Hamiltonian T-manifold with proper moment map ϕ, where T is a connected compact torus. Let c be a value on a singular face of im(ϕ), and let a be a regular value very near c. Then $\pi_1(M_c) \cong \pi_1(M_a)$.*

Lemma 5.4. *(see Lemma 6.15 in* [13]*) Let (M, ω) be a connected Hamiltonian G-manifold with proper moment map ϕ, where G is a connected compact non-abelian Lie group. Let $\tau^P \subset \mathfrak{t}^*_+$ be the principal face. Then, for all value a's on τ^P, $\pi_1(M_a)$'s are isomorphic.*

Lemma 5.5. *(see Lemma 6.19 in* [13]*) Let M be a connected Hamiltonian G-manifold with proper moment map ϕ, where G is a connected compact non-abelian Lie group. Let C be the central face of $\mathfrak{t}^*_+$, and assume that $C \cap$ im$(\phi) \neq \emptyset$ and that C is not the only face of $\mathfrak{t}^*_+$ which contains im(ϕ). Then, for each isotropy type $M_{(H)} \cap \phi^{-1}(C)$, the link of $\left(M_{(H)} \cap \phi^{-1}(C)\right)/G$ in M/G is connected and simply connected.*

We can use Lemma 5.5 to prove Lemma 5.6 below. See the proof of Lemma 6.16 in [13].

Lemma 5.6. *Let (M, ω) be a connected Hamiltonian G-manifold with proper moment map ϕ, where G is a connected compact non-abelian Lie group. Let a be a generic value on the principal face τ^P of $\mathfrak{t}^*_+$, and let c be a value which is not on τ^P. Then $\pi_1(M_c) \cong \pi_1(M_a)$.*

Lemmas 5.1, 5.3, 5.4 and 5.6 imply Proposition 3.

Proposition 3. *Let (M, ω) be a connected Hamiltonian G-manifold with proper moment map ϕ, where G is a connected compact Lie group. Then, $\pi_1(M_a)$'s are isomorphic for all $a \in$ im(ϕ).*

Similar to the proof of Theorem 1.6 in [13], we can use Lemmas 5.2 and 5.5 to prove Proposition 4 which is (A) of Theorem 3.

Proposition 4. *Let* (M, ω) *be a connected Hamiltonian* G*-manifold with proper moment map* ϕ*, where* G *is a connected compact Lie group. Then,* $\pi_1(M/G) \cong \pi_1(M_a)$ *for all* $a \in im(\phi)$*.*

6. Proof of (B) of Theorem 3 for $G = T$

Lemma 6.1. *Let* (M, ω) *be a connected Hamiltonian* T*-manifold with proper moment map* ϕ*, where* T *is a connected compact torus. Let* $T' \subset T$ *be a connected subgroup. Then, for all values* a*'s in one connected chamber of* $im(\phi)$*,* $\pi_1\left(\phi^{-1}(a)/T'\right)$*'s are isomorphic.*

Proof. This is because for all values a's in one connected chamber of im(ϕ), $\phi^{-1}(a)$'s are T-equivariantly diffeomorphic. □

Lemma 6.2. *Assume the assumptions of Theorem 3 hold, where* $G = T$ *is a connected compact torus. Let* $\widehat{T} \subset T$ *be a connected subgroup such that up to a finite subgroup, every isotropy group is a subgroup of* $\widehat{T}$*. Let* $\mathcal{F}$ *be a singular face on* $im(\phi)$*. Let* $\bar{U}$ *be the closure of one open connected chamber* U *such that* $\mathcal{F} \subset \bar{U}$*. Let* N *be* M *or be* $\phi^{-1}(\bar{U})$*. For each isotropy type* $M_H \cap \phi^{-1}(\mathcal{F})$*, let* L_H *be the link of* $\left(M_H \cap \phi^{-1}(\mathcal{F})\right)/T$ *in* N/T*, and let* L'_H *be the link of* $\left(M_H \cap \phi^{-1}(\mathcal{F})\right)/\widehat{T}$ *in* $N/\widehat{T}$*.*

- *If* $H \subseteq \widehat{T}$*, then* $L_H = L'_H$*.*
- *If* $H \subseteq \widehat{T}$ *except for a finite subgroup* Γ *of* H*, then* $L_H = L'_H/\Gamma$*, and,* L_H *and* L'_H *are homeomorphic.*

Proof. Let us prove the claims for $N = M$. Let $p \in M$ be a point with stabilizer H such that $\phi(T \cdot p) \in \mathcal{F}$. By the local normal form theorem (Theorem 6), a neighborhood of the (isotropic) orbit $T \cdot p$ in M is isomorphic to $A = T \times_H (\mathfrak{h}^\circ \times V) = T \times_H (\mathfrak{h}^\circ \times W \times V^H)$, where $\mathfrak{h}$ =Lie(H), $\mathfrak{h}^\circ$ is its annihilator in $\mathfrak{t}^*$, V^H is the subspace of V fixed by H and W is such that $V = W \oplus V^H$. The moment map ϕ on A is $\phi([t, a, w, v]) = \phi(p) + a + \psi(w)$, where ψ is the moment map of the H action on W. The isotropy type $A_H \cap \phi^{-1}(\mathcal{F}) = T \times_H (\mathbb{R}^m \times 0 \times V^H)$, where $\mathbb{R}^m \subset \mathfrak{h}^\circ$ is a subspace which is mapped to $\mathcal{F}$. Let $\mathbb{R}^l$ be the subspace such that $\mathfrak{h}^\circ = \mathbb{R}^l \oplus \mathbb{R}^m$ (note that H acts on $\mathfrak{h}^\circ$ by the trivial coadjoint action).

We see that $\left(A_H \cap \phi^{-1}(\mathcal{F})\right)/T = \mathbb{R}^m \times 0 \times V^H$, and $A/T = \mathfrak{h}^\circ \times V^H \times W/H$. So $L_H = S(\mathbb{R}^l \times W)/H$. Here, $S(\mathbb{R}^l \times W)$ denotes a sphere of the vector space $\mathbb{R}^l \times W$.

If $H \subseteq \widehat{T}$, then $\left(A_H \cap \phi^{-1}(\mathcal{F})\right)/\widehat{T} = T/\widehat{T} \times \mathbb{R}^m \times 0 \times V^H$, and $A/\widehat{T} = T/\widehat{T} \times \mathfrak{h}^\circ \times V^H \times W/H$. So $L'_H = S(\mathbb{R}^l \times W)/H$.

If $H \subseteq \widehat{T}$ except for a finite subgroup Γ, then $\left(A_H \cap \phi^{-1}(\mathcal{F})\right)/\widehat{T} = T/\widehat{T} \times_H \left(\mathbb{R}^m \times 0 \times V^H\right) = T/\widehat{T} \times_\Gamma \left(\mathbb{R}^m \times V^H\right)$, and $A/\widehat{T} = T/\widehat{T} \times_H \left(\mathfrak{h}^\circ \times V^H \times W\right) = (T/\widehat{T} \times \mathbb{R}^m \times V^H) \times_\Gamma \left((\mathbb{R}^l \times W)/(H \cap \widehat{T})\right)$. So $L'_H = S(\mathbb{R}^l \times W)/(H \cap \widehat{T})$. Therefore, $L_H = L'_H/\Gamma = S(\mathbb{R}^l \times W)/(H \cap \widehat{T} \times \Gamma)$.

Since Γ acts on $\mathbb{R}^l$ trivially and it acts on W as a subgroup of $T = (S^1)^m$ (for some m), we easily see that L_H and L'_H are homeomorphic.

We can similarly prove the claims for $\bar{U}$. We refer to the proof of Lemma 3.8 in [13] for the computation of L_H. Then one can compute L'_H as above. □

Lemma 6.3. *Assume the assumptions of Theorem 3 hold, where $G = T$ is a connected compact torus. Let $\widehat{T} \subset T$ be a connected subgroup such that up to a finite subgroup, every isotropy group is a subgroup of $\widehat{T}$. Let $\mathcal{F}$ be a singular face on im(ϕ). Let $\bar{U}$ be the closure of one open connected chamber U such that $\mathcal{F} \subset \bar{U}$.*

1. *For each* nonprincipal isotropy type $M_{(H)} \cap \phi^{-1}(\mathcal{F})$, *the link of $\left(M_{(H)} \cap \phi^{-1}(\mathcal{F})\right)/\widehat{T}$ in $M/\widehat{T}$ is connected and simply connected.*
2. *For each isotropy type $M_{(H)} \cap \phi^{-1}(\mathcal{F})$, the link of $\left(M_{(H)} \cap \phi^{-1}(\mathcal{F})\right)/\widehat{T}$ in $\phi^{-1}(\bar{U})/\widehat{T}$ is connected and simply connected.*

Proof. This follows from Lemmas 5.2 and 6.2. □

Lemma 6.4. *Assume the assumptions of Theorem 3 hold, where $G = T$ is a connected compact torus. Let $\widehat{T} \subset T$ be a connected subgroup such that up to a finite subgroup, every isotropy group is a subgroup of $\widehat{T}$. Let c be a singular value, and let a be a regular value very near c. Let O be a small open neighborhood of c containing a. Let O' be the intersection of O with the connected open chamber containing a, and let $\bar{O}'$ be its closure in O. Let B be the set of values in $\bar{O}'$ but not in O'. Then*

$$\pi_1\left(\phi^{-1}(\bar{O}')/\widehat{T}\right) \cong \pi_1\left(\phi^{-1}(\bar{O}')/\widehat{T} - \phi^{-1}(B)/\widehat{T}\right).$$

Proof. We inductively use (2) of Lemma 6.3 to remove $\phi^{-1}(B)/\widehat{T}$ from $\phi^{-1}(\bar{O}')/\widehat{T}$. One may refer to the proof of Lemma 3.8 in [13]. □

Lemma 6.5. *Assume the assumptions of Theorem 3 hold, where $G = T$ is a connected compact torus. Let c be a singular value, and let a be a regular value very near c. Then $\pi_1\left(\phi^{-1}(c)/\widehat{T}\right) \cong \pi_1\left(\phi^{-1}(a)/\widehat{T}\right)$.*

Proof. We take the O in Lemma 6.4 small enough so that $\phi^{-1}(O)$ and $\phi^{-1}(\bar{O}')$ equivariantly deformation retracts to $\phi^{-1}(c)$ (Theorem 5). So

$$\pi_1\left(\phi^{-1}(\bar{O}')/\widehat{T}\right) \cong \pi_1\left(\phi^{-1}(c)/\widehat{T}\right).$$

Since

$$\pi_1\left(\phi^{-1}(\bar{O}')/\widehat{T} - \phi^{-1}(B)/\widehat{T}\right) \cong \pi_1\left(\phi^{-1}(a)/\widehat{T}\right),$$

the claim follows from Lemma 6.4. □

Lemma 6.6. *Assume the assumptions of Theorem 3 hold, where $G = T$ is a connected compact torus. Let $\widehat{T} \subset T$ be a connected subgroup such that up to a finite subgroup, every isotropy group is a subgroup of $\widehat{T}$. Let*

$\mathcal{F}$ be a singular face on im(ϕ). Let O be a small open neighborhood of $\mathcal{F}$ on the moment map image (O does not intersect the faces which are in the closure of $\mathcal{F}$). Let S be the set of nonprincipal orbits in $\phi^{-1}(\mathcal{F})$. Then $\pi_1\left(\phi^{-1}(O)/\widehat{T}\right) \cong \pi_1\left(\phi^{-1}(O)/\widehat{T} - S/\widehat{T}\right)$.

Proof. The argument is similar to the proof of Lemma 7.20 in [13], but using (1) of Lemma 6.3. $\square$

Proof of (B) of Theorem 3 for $G = T$. Using Lemmas 6.1, 6.5 and 6.6, follow the argument used in the proof of Theorem 1.6 in [13] for the case $G = T$. $\square$

7. PROOF OF (B) OF THEOREM 3 FOR NONABELIAN G

Lemma 7.1. *Assume the assumptions of Theorem 3 hold, where G is nonabelian. Let C be the central face of $\mathfrak{t}^*_+$, and assume that $C \cap im(\phi) \neq \emptyset$ and that C is not the only face of $\mathfrak{t}^*_+$ which contains im(ϕ). For each isotropy type $M_{(H)} \cap \phi^{-1}(C)$, let L_H be the link of $\left(M_{(H)} \cap \phi^{-1}(C)\right)/G$ in M/G, and let L'_H be the link of $\left(M_{(H)} \cap \phi^{-1}(C)\right)/\widehat{G}$ in $M/\widehat{G}$.*

- *If $H \subseteq \widehat{G}$, then $L_H = L'_H$, they are connected and $\pi_1(L_H) = \pi_1(L'_H) = 1$.*
- *If $H \subseteq \widehat{G}$ except for finitely many elements, then L_H and L'_H are connected and $\pi_1(L_H) = \pi_1(L'_H) = 1$.*

Proof. By the local normal form theorem (Theorem 6), a neighborhood in M of an orbit O in $\phi^{-1}(C)$ with isotropy group (H) is isomorphic to $A = G \times_H (\mathfrak{h}^\circ \times V) = G \times_H (\mathfrak{h}^\circ \times W \times V^H)$, where $\mathfrak{h}^\circ$ is the annihilator of $\mathfrak{h}$ =Lie(H) in $\mathfrak{g}^*$ =Lie*(G), and, V^H is the subspace of V fixed by H and W is such that $V = W \oplus V^H$. The moment map ϕ on A is $\phi([g, a, w, v]) = Ad^*(g)\,(\phi(\mathrm{O}) + a + \psi(w))$, where ψ is the moment map of the H action on W. So $A_{(H)} \cap \phi^{-1}(C) = G \times_H (\mathbb{R}^m \times 0 \times V^H)$, where $\mathbb{R}^m \subset \mathfrak{h}^\circ$ is the subspace which is mapped to C and therefore on which H acts trivially. Let $\mathbb{R}^l \subset \mathfrak{h}^\circ$ be the subspace such that $\mathfrak{h}^\circ = \mathbb{R}^l \oplus \mathbb{R}^m$.

Then $\left(A_{(H)} \cap \phi^{-1}(C)\right)/G = \mathbb{R}^m \times 0 \times V^H$, and $A/G = \mathbb{R}^m \times 0 \times V^H \times (\mathbb{R}^l \times W)/H$. So $L_H = S(\mathbb{R}^l \times W)/H$.

If $H \subseteq \widehat{G}$, then $\left(A_{(H)} \cap \phi^{-1}(C)\right)/\widehat{G} = G/\widehat{G} \times \mathbb{R}^m \times V^H$, and $A/\widehat{G} = G/\widehat{G} \times \mathbb{R}^m \times V^H \times (\mathbb{R}^l \times W)/H$. So $L'_H = S(\mathbb{R}^l \times W)/H = L_H$.

If $H \subseteq \widehat{G}$ except for finitely many elements of H, then $\left(A_{(H)} \cap \phi^{-1}(C)\right)/\widehat{G}$ $= G/\widehat{G} \times_H (\mathbb{R}^m \times V^H) = (\mathbb{R}^m \times V^H) \times G/\widehat{G} \times_H 0$, and $A/\widehat{G} = G/\widehat{G} \times_H (\mathbb{R}^m \times V^H \times \mathbb{R}^l \times W) = (\mathbb{R}^m \times V^H) \times G/\widehat{G} \times_H (\mathbb{R}^l \times W)$. So $L'_H = S(\mathbb{R}^l \times W)/(H \cap \widehat{G})$.

By Lemma 5.5, L_H is connected and $\pi_1(L_H) = 1$. The claim for the first case when $L_H = L'_H$ follows immediately. For the second case, we think about it as follows. If we take the new subgroup $H' = H \cap \widehat{G}$ of G which differs from H by finitely many elements, the $S(\mathbb{R}^l \times W)$ in the above analysis does not change; by the same arguments (or criteria) of the connectedness and simply connectedness of L_H, L'_H is connected and simply connected. For details of the arguments, we refer to the proof of Lemma 6.19 in [13]. $\square$

Lemma 7.2. *Assume the assumptions of Theorem 3 hold, where G is non-abelian. Let C be the central face of $\mathfrak{t}^*_+$, and assume that $C \cap im(\phi) \neq \emptyset$ and that C is not the only face of $\mathfrak{t}^*_+$ which contains $im(\phi)$. Then*

$$\pi_1\left(M/\widehat{G}\right) \cong \pi_1\left(M/\widehat{G} - \phi^{-1}(C)/\widehat{G}\right).$$

Similarly, if O is a small open invariant neighborhood of C in $\mathfrak{g}^*$, then

$$\pi_1\left(\phi^{-1}(O)/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(O)/\widehat{G} - \phi^{-1}(C)/\widehat{G}\right).$$

Proof. By Lemma 7.1, for each possible H, the link of $\left(M_{(H)} \cap \phi^{-1}(C)\right)/\widehat{G}$ in $M/\widehat{G}$ is connected and simply connected. Therefore we can inductively remove the strata of $\phi^{-1}(C)/\widehat{G}$ from $M/\widehat{G}$ or from $\phi^{-1}(O)/\widehat{G}$ using Lemma 2.2. $\square$

Lemma 7.3. *Assume the assumptions of Theorem 3 hold, where G is non-abelian.*

(1) *Let τ be a non-principal face of $\mathfrak{t}^*_+$ such that $\tau \cap im(\phi) \neq \emptyset$ and let R^τ be the corresponding cross section. Then, for each $M_{(H)} \cap \phi^{-1}(\tau)$, the link of $\left(M_{(H)} \cap \phi^{-1}(\tau)\right)/(G_\tau \cap \widehat{G})$ in $R^\tau/(G_\tau \cap \widehat{G})$ is the same as the link of $\left(M_{(H)} \cap \phi^{-1}(G \cdot \tau)\right)/\widehat{G}$ in $G \cdot R^\tau/\widehat{G}$.*

(2) *Let τ^P be the principal face and let R^{τ^P} be the principal cross section. Let $c \in \tau^P$ be a singular value. Then, for each $M_{(H)} \cap \phi^{-1}(c)$, the link of $\left(M_{(H)} \cap \phi^{-1}(c)\right)/(T \cap \widehat{G})$ in $R^{\tau^P}/(T \cap \widehat{G})$ is the same as the link of $\left(M_{(H)} \cap \phi^{-1}(G \cdot c)\right)/\widehat{G}$ in $G \cdot R^{\tau^P}/\widehat{G}$.*

Proof. First note that, if a point with isotropy group H is mapped to a face τ, then since $\phi(g \cdot m) = Ad^*(g) \cdot \phi(m)$, $H \subset G_\tau$.

(1) The manifold $G \cdot R^\tau$ is a fibration over the coadjoint orbit G/G_τ with fiber R^τ; and correspondingly, each $M_{(H)} \cap (G \cdot R^\tau)$ is a fibration over G/G_τ with fiber $M_{(H)} \cap R^\tau$, moreover, each $M_{(H)} \cap \phi^{-1}(G \cdot \tau)$ is a fibration over G/G_τ with fiber $M_{(H)} \cap \phi^{-1}(\tau)$. The claim follows from the equivariance of ϕ.

The proof of (2) is similar. $\square$

Remark 7.4. Note that for any face $\tau \subset \mathfrak{t}^*_+$ and the stabilizer group G_τ of τ, the subgroup $G_\tau \cap \widehat{G}$ of G_τ is connected since G_τ and $\widehat{G}$ are both connected.

Lemma 7.5. *Assume the assumptions of Theorem 3 hold, where G is non-abelian. Let $\tau^P \subset \mathfrak{t}^*_+$ be the principal face. Then, $\pi_1\left(\phi^{-1}(G \cdot \tau^P)/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(G \cdot a)/\widehat{G}\right)$ for all $a \in \tau^P$.*

Proof. First, consider the space $\phi^{-1}(\tau^P)$ with the T action, where T is a maximal torus of G. Similar to the proof of the theorem for abelian group actions, by using removing and deforming in the space $\phi^{-1}(\tau^P)/(\widehat{G}\cap T)$, we get

$$\pi_1\left(\phi^{-1}(\tau^P)/(\widehat{G}\cap T)\right) \cong \pi_1\left(\phi^{-1}(a)/(\widehat{G}\cap T)\right)$$

for some *particular* $a \in \tau^P$. Then by Lemma 6.3 and (2) of Lemma 7.3, we can do corresponding removing and deforming in the space $\phi^{-1}(G\cdot\tau^P)/\widehat{G}$, and we arrive at

$$\pi_1\left(\phi^{-1}(G\cdot\tau^P)/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(G\cdot a)/\widehat{G}\right)$$

for this *particular* $a \in \tau^P$.

For any two values a and b in the same connected open face on τ^P,

$$\pi_1\left(\phi^{-1}(G\cdot a)/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(G\cdot b)/\widehat{G}\right)$$

since $\phi^{-1}(G\cdot a)$ and $\phi^{-1}(G\cdot b)$ are equivariantly diffeomorphic. Now, let $c \in \tau^P$ be a value on a singular face. Take a value $a \in \tau^P$ in a connected open face very close to c. Take $\bar{O}' \subset \tau^P$ as in Lemma 6.4. Using (2) of Lemma 7.3 again, we can do removing in $\phi^{-1}(G\cdot\bar{O}')/\widehat{G}$ corresponding to the removing in $\phi^{-1}(\bar{O}')/(T\cap\widehat{G})$; and we arrive at

$$\pi_1\left(\phi^{-1}(G\cdot a)/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(G\cdot c)/\widehat{G}\right).$$

□

Lemma 7.6. *Assume the assumptions of Theorem 3 hold, where* G *is non-abelian. Let* $c \in \tau$ *be a value, where* τ *is a face of* $\mathfrak{t}^*_+$ *such that* $\tau \neq \tau^P$*, and let* a *be a generic value on* τ^P *very near* c*. Let* $O \subset \mathfrak{g}^*$ *be a small open invariant neighborhood of* c *containing* a*. Let* B *be the set of values in* $O \cap \mathfrak{t}^*_+$ *other than those on the open connected face of generic values on* τ^P *containing* a*. Then*

$$\pi_1\left(\phi^{-1}(O)/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(O)/\widehat{G} - \phi^{-1}(G\cdot B)/\widehat{G}\right).$$

Proof. Consider the cross section R^τ where G_τ acts. Note that τ lies in the central dual Lie algebra of G_τ. Using Lemmas 7.2 and 7.3, we have

$$\pi_1\left(\phi^{-1}(O)/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(O)/\widehat{G} - \phi^{-1}(G\cdot\tau)/\widehat{G}\right).$$

For other non-principal faces τ''s, we use the cross section theorem and Lemma 7.3 to inductively remove $\phi^{-1}(G\cdot\tau')/\widehat{G}$'s. If there are singular faces on $O\cap\tau^P$, then we use Lemmas 6.3 and 7.3 to remove the rest (deforming may also be needed). If further detail is prefered, one may refer to the proof of Lemma 6.18 in [13]. □

Lemma 7.7. *Assume the assumptions of Theorem 3 hold, where G is nonabelian. Let $c \in \tau$ be a value, where τ is a face of $\mathfrak{t}^*_+$ such that $\tau \neq \tau^P$, and let a be a generic value on τ^P very near c. Then* $\pi_1\left(\phi^{-1}(G\cdot c)/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(G\cdot a)/\widehat{G}\right)$.

Proof. In Lemma 7.6, we take O small enough such that $\phi^{-1}(O)$ equivariantly deformation retracts to $\phi^{-1}(G\cdot c)$. Then

$$\pi_1\left(\phi^{-1}(O)/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(G\cdot c)/\widehat{G}\right).$$

Since

$$\pi_1\left(\phi^{-1}(O)/\widehat{G} - \phi^{-1}(G\cdot B)/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(G\cdot a)/\widehat{G}\right),$$

the claim follows from Lemma 7.6. □

Proof of (B) of Theorem 3 for nonabelian G. We use Lemma 7.2, the cross section theorem and Lemma 7.3 to inductively remove $\phi^{-1}(G\cdot\tau)/\widehat{G}$ from $M/\widehat{G}$ for the faces τ's other than the principal face τ^P which contain values of ϕ. Assume we have now $\pi_1(M/\widehat{G}) \cong \pi_1\left(\phi^{-1}(G\cdot\tau^P)/\widehat{G}\right)$. Then by Lemma 7.5, $\pi_1\left(\phi^{-1}(G\cdot\tau^P)/\widehat{G}\right) \cong \pi_1\left(\phi^{-1}(G\cdot a)/\widehat{G}\right)$ for $a \in \tau^P$. The claim that the $\pi_1\left(\phi^{-1}(G\cdot a)/\widehat{G}\right)$'s are isomorphic for all $a \in \mathrm{im}(\phi)$ follows from Lemma 7.7. □

REFERENCES

[1] M. Atiyah, *Convexity and commuting Hamiltonians*, Bull. Lond. Math. Soc. **14** (1982), 1-15.

[2] G. Bredon, *Introduction to Compact Transformation Groups*, Academic Press, New York, 1972.

[3] V. Guillemin, E. Lerman, and S. Sternberg, *Symplectic fibrations and multiplicity diagrams*, Cambridge University Press, Cambridge, 1996.

[4] V. Guillemin and S. Sternberg, *Convexity properties of the moment mapping*, Invent. Math. **67** (1982), no. 3, 491–513.

[5] V. Guillemin and S. Sternberg, *Symplectic techniques in physics*, Cambridge University Press, Cambridge, 1990.

[6] V. Guillemin and S. Sternberg, *A normal form for the moment map*, Differential geometric methods in mathematical physics, (S. Sternberg, Ed.) Reidel, Dordrecht, Holland, 1984.

[7] L. Godinho and M. E. Sousa-Dias, *The fundamental group of S^1-manifolds*, math.SG/0505420.

[8] F. C. Kirwan, *Convexity properties of the moment mapping III*, Invent. Math. **77** (1984), 547-552.

[9] F. C. Kirwan, *The cohomology of quotients in symplectic and algebraic geometry*, Princeton University Press, 1984.

[10] E. Lerman, *Gradient flow of the norm squared of a moment map*, Enseign.Math. (2) 51 (2005), no. 1-2, 117-127.

[11] E. Lerman, E. Meinrenken, S. Tolman, and C. Woodward, *Nonabelian convexity by symplectic cuts*, Topology **37**, no. 2, 245–259, 1998.

[12] H. Li, *π_1 of Hamiltonian S^1-manifolds*, Proc. Amer. Math. Soc. **131**, no. 11, 3579-3582, 2003.

[13] H. Li, *The fundamental group of symplectic manifolds with Hamiltonian Lie group actions*, Journal of Symplectic Geometry, Vol. 4, No. 3, 345-372, 2007.

[14] C. M. Marle, *Le voisinage d'une orbite d'une action hamiltonienne d'un group de Lie*, In: Séminaire sud-rhodanien de géométrie, Π (Lyon, 1983), 19-35, Travaux en cours, Hermann, Paris, 1984.

[15] R. Sjamaar and E. Lerman, *Stratified symplectic spaces and reduction*, Ann. of Math. (2) **134**, no. 2, 375–422, 1991.

[16] C. Woodward, *The Yang-Mills heat flow on the moduli space of framed bundles on a surface*, Amer. J. Math. 128 (2006), no. 2, 311-359.

Institut de Mathematiques de Bourgogne, 9 Av. Alain Savary, 21078, Dijon, France.

E-mail address: `hui.li@u-bourgogne.fr`